

Serial No. 10/751,009
Docket No. SHE0074.00

In the Claims:

Please amend claim 35 as indicated below. Currently amended claims are presented with markings to indicate the changes made, wherein a ~~strike through~~ is used to designate deletions and underlining is used to designate additions.


RECEIVED
CENTRAL FAX CENTER

SEP 28 2006

Claims 1-34 (Canceled).

35. (Currently Amended) A polymeric reagent comprising a water-soluble polymer segment and a functional group, wherein the functional group is part of a cyclic structure and further wherein the cyclic structure is attached to the water-soluble polymer segment through either a direct covalent bond or through one or more atoms, wherein the functional group is selected from the group consisting of ketone, ketone hydrate, thione, monothiohydrate, dithiohydrate, hemiketal, monothiohemiketal, dithiohemiketal, ketal, and dithioketal, and further wherein (i) the polymeric reagent lacks a cyclic dienone, and (ii) a carbon atom is attached on each side of the functional group.

36. (Original) The polymeric reagent of claim 35, comprising the following structure:




wherein:

- POLY¹ is a water-soluble polymer segment having a terminus defined as -O-R⁷;
- (a) is either zero or one;
- (b) is either zero or one;
- X¹, when present, is a spacer moiety;
- EW¹, when present, is an electron-withdrawing group;
- (z) is zero or a positive an integer;
- each occurrence of R², when present, is independently H or an organic radical;
- each occurrence R³, when present, is independently H or an organic radical; and
- R⁷ is H or an organic radical.

Serial No. 10/751,009
Docket No. SHE0074.00

37. (Original) The polymeric reagent of claim 35, comprising the following structure:



wherein:

- POLY¹ is a water-soluble polymer segment having a terminus defined as -O-R⁷;
- (a) is either zero or one;
- (b) is either zero or one;
- X¹, when present, is a spacer moiety;
- EW¹, when present, is an electron-withdrawing group;
- (z) is zero or a positive an integer;
- each occurrence of R², when present, is independently H or an organic radical;
- each occurrence R³, when present, is independently H or an organic radical; and
- R⁷ is H or an organic radical.

38. (Original) The polymeric reagent of claims 36 or 37, wherein the water-soluble polymer segment ("POLY") is a poly(ethylene glycol).

39. (Previously Presented) The polymeric reagent of claim 38, wherein the poly(ethylene glycol) is -CH₂CH₂-(OCH₂CH₂)_m-OCH₂CH₂-, wherein (m) is from about 3 to about 3,000.

40. (Original) The polymeric reagent of claim 38, wherein the poly(ethylene glycol) is branched.

41. (Original) The polymeric reagent of claims 36 or 37, wherein R⁷ is a nonaromatic organic radical comprising nine or less carbon atoms.

42. (Original) The polymeric reagent of claims 36 or 37, wherein R⁷ is H or methyl.


Serial No. 10/751,009
Docket No. SHE0074.00

43. (Original) The polymeric reagent of claim 38, wherein the poly(ethylene glycol) has a nominal average molecular weight of from about 2,500 Daltons to about 100,000 Daltons.
44. (Original) The polymeric reagent of claim 43, wherein the poly(ethylene glycol) has a nominal average molecular weight of from about 4,900 Daltons to about 40,000 Daltons.
45. (Original) The polymeric reagent of claim 44, wherein the poly(ethylene glycol) has a nominal average molecular weight of from about 9,900 Daltons to about 25,000 Daltons.
46. (Original) The polymeric reagent of claim 35, wherein the functional group is selected from the group consisting of thione, monothiohydrate, dithiohydrate, monothiohemiketal, dithiohemiketal, and dithioketal.
47. (Original) The polymeric reagent of claim 35, lacking halogen atoms.
48. (Original) The polymeric reagent of claims 36 or 37, lacking an aromatic moiety.
49. (Original) The polymeric reagent of claims 36 or 37, lacking an enone.
50. (Original) The polymeric reagent of claims 36 or 37, wherein (z) is zero, (b) is one, and EW¹ is selected from the group consisting of -O-, -NH-, -NHC(O)-, -C(O)NH-, -OC(O)-, -OC(O)-, -OC(O)-NH-, -NH-OC(O)-, -C(O)-, -C(S)-, and -C(OR)H-, wherein OR is an alkoxy or hydroxy substituent.
51. (Original) The polymeric reagent of claim 50, wherein EW¹ is -C(O)-.
52. (Original) The polymeric reagent of claim 35, wherein the cyclic structure is selected from the group consisting of piperidinyl, cyclohexyl, cyclopentyl, bicyclo[2,2,1]heptanyl, bicyclo[2,2,1]hexanyl, bicyclo[3,2,1]octanyl, bicyclo[3,1,1]heptanyl,

Serial No. 10/751,009
Docket No. SHE0074.00

bicyclo[3,3,1]nonanyl, cyclobutyl, cycloheptyl, cyclooctyl, oxa and aza forms thereof, and dioxo and diaza forms of any of the foregoing.

53. (Previously Presented) The polymeric reagent of claim 35, selected from the group consisting of:



wherein (m) is from about 3 to about 3000.

Claims 54-169 (Canceled).

170. (Withdrawn) A method for making a polymer-active agent conjugate comprising contacting, under conjugation conditions, an active agent with a polymeric reagent comprised of a water-soluble polymer segment attached, either directly or through one or more atoms, to a functional group, wherein the functional group is selected from the group consisting of ketone, ketone hydrate, thione, monothiohydrate, dithiohydrate, hemiketal, monothiohemiketal, dithiohemiketal, ketal, and dithioketal.

171. (Withdrawn) The method of claim 170, wherein the polymeric reagent is encompassed by claim 35.

172. (Withdrawn) The method of claim 170, wherein the functional group s a ketone.

Serial No. 10/751,009
Docket No. SHE0074.00

173. (Withdrawn) The method of claim 170, wherein the functional group is a ketone hydrate.
174. (Withdrawn) The method of claim 170, wherein the functional group is a hemiketal.
175. (Withdrawn) The method of claim 170, wherein the functional group is a ketal.
176. (Withdrawn) The method of claim 170, wherein the functional group is a thione.
177. (Withdrawn) The method of claim 170, wherein the functional group is a monothiohydrate.
178. (Withdrawn) The method of claim 170, wherein the functional group is a dithiohydrate.
179. (Withdrawn) The method of claim 170, wherein the functional group is a monothiohemiketal,
180. (Withdrawn) The method of claim 170, wherein the functional group is a dithiohemiketal.
181. (Withdrawn) The method of claim 170, wherein the functional group is a dithioketal.
182. (Withdrawn) The method of claim 170, wherein the active agent is a polypeptide.
183. (Canceled).
184. (Withdrawn) The method of claim 170, wherein the active agent is not a polypeptide.

Serial No. 10/751,009
Docket No. SHE0074.00

185. (Canceled).

186. (Canceled).

187. (Withdrawn) A conjugate prepared according to the method of claim 170.

188. (Withdrawn) A pharmaceutical preparation comprising the conjugate of claim 187 and a pharmaceutically acceptable excipient.

(This space intentionally left blank.)